

Safe sex practices/ STI prevention

Skin cancer prevention

Adult Health Maintenance Form

| Name: | | | |
|-------|--|--|--|
| | | | |

MR #:

| Preventive Services (Record results, notes, etc. in box) | | Date / / | Date / / | Date / / |
|--|--|---|--|-------------------------------------|
| Well-person exam | | | | |
| Alcohol & recreational drug assessment (See reverse side for Cenpatico referral info) | | □ Cenpatico referral | □ Cenpatico referral | □ Cenpatico referral |
| Blood pressure | | | | |
| Body Mass Index weight(kg)/ height (m |)2 | | | |
| Cholesterol screening (Note collection date and result) | | _/_/ _ LDL: | _/_/ _ LDL: | _/_/ _ LDL: |
| Colorectal screening (Please note if FOBT/colonoscopy, date and result) | | □NA | □NA | □NA |
| Depression scree (See reverse side for Ph | ning/ PHQ – 2 IQ-2 and Cenpatico referral info) | □ Cenpatico referral | □ Cenpatico referral | □ Cenpatico referral |
| | Hemoglobin A1c (Note collection date and result) | _/_/ | | _/_/ |
| Diabetes (If already diagnosed | Urine microalbumin test (Note collection date and result) | | _/_/ | _!_! |
| with diabetes) | Retinal eye exam by an eye care professional (Note exam date and result) | /_/_ □ No retinopathy □ + Retinopathy | /_/ □ No retinopathy □ + Retinopathy | // □ No retinopathy □ + Retinopathy |
| Diabetes Type II screening | | □NA | □NA | □NA |
| Fall risk assessme | ent | □NA | □NA | □NA |
| Hepatitis C screening | | □NA | □NA | □NA |
| HIV screening | | □NA | □NA | □NA |
| Influenza vaccine | | | | |
| Pneumococcal vaccine | | □NA | □NA | □NA |
| Tobacco use assessment (If positive for tobacco use, please consider an Indiana Tobacco Quit Line referral: phone # listed to right) | | □ 1-800-QUIT NOW | □ 1-800-QUIT NOW | □ 1-800-QUIT NOW |
| ♀ Breast cancer s | | □NA | □NA | □NA |
| ☐ Cervical cancer screening (Note collection date and result) | | □NA | □NA | □NA |
| ♀ Chlamydia screening | | □NA | □NA | □NA |
| ♀ Osteoporosis screening | | □NA | □NA | □NA |
| ♂ Abdominal aortic aneurism screening | | □NA | □NA | □NA |
| | | | | |
| Education & Counseling | | (Same date as above) | (Same date as above) | (Same date as above) |
| Advance directive/ Health care proxy | | □ Copy on file | □ Copy on file | □ Copy on file |
| Aspirin chemopro | - | | | |
| Dental exam referral/ Oral hygiene Domestic violence/ Elder abuse | | | | |
| | EILEI ADUSE | | | |
| Fall prevention | | | | |

PHQ- 2: Screening Instrument for Depression

| Over the past two weeks, how often have you been bothered by any of the following problems? | Not at all | Several days | More than one- half the days | Nearly everyday |
|---|------------|--------------|---------------------------------|--------------------|
| Little interest or pleasure in doing things | 0 | 1 | 2 | 3 |
| Feeling down, depressed, or hopeless | 0 | 1 | 2 | 3 |

Note: If the patient has a positive response to either question, consider administering the PHQ-9 or asking the patient more questions about possible depression. For older adults, consider the PHQ-9 or the 15-item Geriatric Depression Scale. A negative response to both questions is considered a negative result for depression. Cenpatico (CBH) is MHS' sister company and the behavioral health vendor available to all MHS members. PMPs, at any time, can refer members for case management services who may benefit from coordination of activities or additional assistance in obtaining resources. Please contact Cenpatico at 1-877-647-4848 to make a referral or to collaborate about a member's care. A referral form to Cenpatico's Disease Management program is also available at mhsindiana.com/for-providers/provider-forms/

| Recommended Services | Frequency | Recommended Services | Frequency |
|-------------------------------------|--|---------------------------------------|--|
| Well-person exam | Annually | Hepatitis C screening | Persons at high risk for infection and adults born between 1945 and 1965 |
| ETOH & recreational drug assessment | Annually | HIV screening | All adolescents and adults ages 15 to 65 years and others who are at increased risk for HIV infection and all pregnant women |
| Aspirin chemoprophylaxis | When the potential harm of an increase in gastrointestinal hemorrhage is outweighed by a potential benefit of a reduction in myocardial infarctions (men aged 45-79 years) or in ischemic strokes (women aged 55-79 years). | Influenza vaccine | Annually |
| Blood pressure | -Screening every 2 years with BP <120/80Screening every year with SBP of 120-139 mmHg or DBP of 80-90 mmHg | Pneumococcal vaccine | Age 19 and older- those at increased risk All adults age 65 and older |
| Body Mass Index | Patients with a body mass index of 30 kg/m2 or higher should be offered or referred to intensive, multicomponent behavioral interventions | Tobacco use assessment | Ask all adults about tobacco use and provide tobacco cessation interventions for those who use tobacco; provide augmented, pregnancy-tailored counseling for those pregnant women who smoke |
| Cholesterol screening | Men aged 20-35 and women over age 20 who are at increased risk for coronary heart disease; all men aged 35 and older | ♀ Breast cancer screening | Biennial screening mammography for women aged 50 to 74 years. Note: The Department of Health and Human Services, in implementing the ACA, follows the 2002 USPSTF recommendation for screening mammography, with or without clinical breast examination, every 1-2 years for women aged 40 and older |
| Colorectal screening | Adults aged 50-75 using fecal occult blood testing, sigmoidoscopy, or colonoscopy | ♀ Cervical cancer screening | Screen with cytology every 3 years (women ages 21 to 65) or co-test (cytology/HPV testing) every 5 years (women ages 30-65). |
| Depression screening | Non- pregnant adults 18 years and older. The American Academy of Pediatrics (AAP) recommends children age 11 years and older be screened annually. Recommended screening tools, such as the PHQ-9, are available in the GLAD-PC toolkit. | ♀ Chlamydia screening | Sexually active women 24 and younger and other asymptomatic women at increased risk for infection. Asymptomatic pregnant women 24 and younger and others at increased risk. |
| Diabetes Type II screening | Asymptomatic adults with sustained blood pressure greater than 135/80 mg Hg. | ♀ Osteoporosis screening | Women aged 65 years and older and women under age 65 whose 10-year fracture risk is equal to or greater than that of a 65-year-old white woman without additional risk factors. |
| Fall risk assessment | Provide intervention (exercise or physical therapy and/or vitamin D) to adults ≥65 years at increased risk for falls. | ∂ Abdominal aortic aneurism screening | One-time screening by ultrasound in men aged 65 to 75 who have ever smoked. |

Preventive services guidelines referenced from the US Preventive Services Task Force. For complete guidelines please visit <u>uspreventiveservicestaskforce.org.</u> MHS Clinical Practice Guidelines can be found at <u>mhsindiana.com/for-providers/practice-guidelines/</u>