

# FUH: Follow-Up After Hospitalization for Mental Illness



## Members 6 years of age and older

Who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health practitioner. Two rates are reported:

- The percentage of discharges for which the member received follow-up within 7 days of discharge. Codes to identify follow-up visits (must be with mental health practitioner)
- The percentage of discharges for which the member received follow-up within 30 days of discharge.

Description	CPT	CPT Modifier	HCPCS	POS
An outpatient visit (Visit Setting Unspecified Value Set with Outpatient POS Value Set with a mental health practitioner, with or without a telehealth modifier (Telehealth Modifier Value Set)).	90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 - 99223, 99231 - 99233, 99238, 99239, 99251 - 99255	95, GT	N/A	03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
An outpatient visit (BH Outpatient Value Set with a mental health practitioner, with or without a telehealth modifier (Telehealth Modifier Value Set)).	98960 - 98962, 99078, 99201 - 99205, 99211 - 99215, 99241 - 99245, 99341 - 99345, 99347 - 99350, 99381 - 99387, 99391 - 99397, 99401 - 99404, 99411, 99412, 99510, 99483	95, GT	G0155, G0176, G0177, G0409, G0463, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, M0064, T1015	N/A
An intensive outpatient encounter or partial hospitalization (Visit Setting Unspecified Value Set with Partial Hospitalization POS Value Set with a mental health practitioner, with or without a telehealth modifier (Telehealth Modifier Value Set)).	90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 - 99223, 99231 - 99233, 99238, 99239, 99251 - 99255	95, GT	N/A	52

Description	CPT	CPT Modifier	HCPCS	POS
An intensive outpatient encounter or partial hospitalization (Partial Hospitalization/ Intensive Outpatient Value Set) with a mental health practitioner.	N/A	N/A	G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485	N/A
A community mental health center visit (Visit Setting Unspecified Value Set with Community Mental Health Center POS Value Set with a mental health practitioner, with or without a telehealth modifier (Telehealth Modifier Value Set).	90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 - 99223, 99231 - 99233, 99238, 99239, 99251 - 99255	95, GT		53
Electroconvulsive therapy (Electroconvulsive Therapy Value Set) with (Ambulatory Surgical Center POS Value Set; Community Mental Health Center POS Value Set; Outpatient POS Value Set; Partial Hospitalization POS Value Set) with a mental health practitioner.	90870	N/A	N/A	24, 53, 52, 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
A telehealth visit: Visit Setting Unspecified Value Set with Telehealth POS Value Set with a mental health practitioner, with or without a telehealth modifier (Telehealth Modifier Value Set).	90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 - 99223, 99231 - 99233, 99238, 99239, 99251 - 99255	95, GT	N/A	02
An observation visit (Observation Value Set) with a mental health practitioner.	99217 - 99220	N/A	N/A	N/A
Transitional care management services (Transitional Care Management Services Value Set), with a mental health practitioner, with or without a telehealth modifier (Telehealth Modifier Value Set).	99495, 99496	N/A	N/A	N/A

**To improve HEDIS scores:**

- Ensure members 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses receive the appropriate follow-up visit with a mental health practitioner with 7 day or 30 days of discharge.
- Ensure appropriate coding to capture services provided within the appropriate timeframe..

**IET- INITIATION AND ENGAGEMENT OF ALCOHOL AND OTHER DRUG ABUSE OR DEPENDENT TREATMENT**

Measure evaluates the percentage of adolescent and adult members with a new episode of alcohol or other drug abuse or dependence (AOD) who received:

- Initiated dependence treatment within 14 days of their diagnosis
- Continued treatment with 2 or more additional services within 34 days of the initiation visit

For the follow up treatments, include an ICD-10 diagnosis for Alcohol or Other Drug Dependence from the Mental, Behavioral and Neurodevelopmental Disorder Section of ICD-10 along with a procedure code for the preventive service, evaluation and management consultation or counseling service (see codes below).

**TREATMENT CODES TO BE USED WITH DIAGNOSIS CODES**

CPT	CPT Modifier	HCPCS	POS
98960 - 98962, 99078, 99201 - 99205, 99211 - 99215, 99241 - 99245, 99341 - 99345, 99347 - 99350, 99384 - 99387, 99394 - 99397, 99401 - 99404, 99408 - 99409, 99411 - 99412, 99510, 98969, 99444, 98966 - 98968, 99441 - 99443, 90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875 - 90876, 99221 - 99223, 99231 - 99233, 99238, 99239, 99251 - 99255, 99483	95, GT	G0155, G0176, G0177, G0396, G0397, G0409-G0411, G0443, G0463, H0001, H0002, H0004, H0005, H0007, H0015, H0016, H0022, H0031, H003-H0037, H0039, H0040, H0047, H2000, H2001, H2010-H2020, H2035, H2036, M0064, S0201, S9480, S9484, S9485, T1006, T1012, T1015	02, 03, 05, 07, 09, 11-20, 22, 33, 49- 50, 52-53, 57, 71-72

**To improve HEDIS scores:**

- Ensure adolescent and adult members with a new episode of alcohol or other drug abuse or dependence receive initial dependence treatment within 14 days of diagnosis.
- Ensure members remain adherent to treatment by scheduling 2 or more additional services within 34 days of the initiation visit.
- Ensure appropriate coding to capture services provided within the appropriate timeframe.

## AMM- ANTIDEPRESSANT MEDICATION MANAGEMENT

Measure evaluates the percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment. Two rates are reported.

- Effective Acute Phase Treatment. The percentage of members who remained on an antidepressant medication for at least 84 days (12 weeks).
- Effective Continuation Phase Treatment. The percentage of members who remained on an antidepressant medication for at least 180 days (6 months).
- For the follow up treatments, include an ICD-10 diagnosis for Alcohol or Other Drug Dependence from the Mental, Behavioral and Neurodevelopmental Disorder Section of ICD-10 along with a procedure code for the preventive service, evaluation and management consultation or counseling service (see codes below).

## ANTIDEPRESSANT MEDICATIONS

Description	Prescription
Miscellaneous antidepressants	<ul style="list-style-type: none"> <li>• Bupropion</li> <li>• Vilazodone</li> <li>• Vortioxetine</li> </ul>
Monoamine oxidase inhibitors	<ul style="list-style-type: none"> <li>• Isocarboxazid</li> <li>• Selegiline</li> <li>• Phenelzine</li> <li>• Tranylcypromine</li> </ul>
Phenylpiperazine antidepressants	<ul style="list-style-type: none"> <li>• Nefazodone</li> <li>• Trazodone</li> </ul>
Psychotherapeutic combinations	<ul style="list-style-type: none"> <li>• Amitriptyline-chlordiazepoxide</li> <li>• Amitriptyline-perphenazine</li> <li>• Fluoxetine-olanzapine</li> </ul>
SNRI antidepressants	<ul style="list-style-type: none"> <li>• Desvenlafaxine</li> <li>• Levomilnacipran</li> <li>• Duloxetine</li> <li>• Venlafaxine</li> </ul>
SSRI antidepressants	<ul style="list-style-type: none"> <li>• Citalopram</li> <li>• Fluoxetine</li> <li>• Paroxetine</li> <li>• Escitalopram</li> <li>• Fluvoxamine</li> <li>• Sertraline</li> </ul>
Tetracyclic antidepressants	<ul style="list-style-type: none"> <li>• Maprotiline</li> <li>• Mirtazapine</li> </ul>
Tricyclic antidepressants	<ul style="list-style-type: none"> <li>• Amitriptyline</li> <li>• Desipramine</li> <li>• Nortriptyline</li> <li>• Amoxapine</li> <li>• Doxepin (&gt;6 mg)</li> <li>• Protriptyline</li> <li>• Clomipramine</li> <li>• Imipramine</li> <li>• Trimipramine</li> </ul>

### To improve HEDIS scores:

- Ensure members remain adherent to antidepressant medication treatment. Ongoing monitoring is critical to adherence.