

Clinical Policy: Entrectinib (Rozlytrek)

Reference Number: CP.PHAR.441

Effective Date: 12.01.19 Last Review Date: 11.20

Line of Business: Commercial, HIM, Medicaid Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Entrectinib (Rozlytrek[™]) is a kinase inhibitor.

FDA Approved Indication(s)

Rozlytrek is indicated for the treatment of:

- Adult patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are ROS1-positive.
- Adult and pediatric patients 12 years of age and older with solid tumors that:
 - o have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation,
 - o are metastatic or where surgical resection is likely to result in severe morbidity, and
 - o have either progressed following treatment or have no satisfactory alternative therapy.

This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Rozlytrek is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Non-Small Cell Lung Cancer (must meet all):
 - 1. Diagnosis of recurrent, advanced, or metastatic NSCLC;
 - 2. Prescribed by or in consultation with an oncologist;
 - 3. Age \geq 18 years;
 - 4. Disease is ROS1 positive;
 - 5. Member has not received prior ROS1 targeted therapy (e.g., Xalkori[®], Zykadia[®], Lorbrena[®]);
 - 6. Request meets one of the following (a or b):*
 - a. Dose does not exceed 600 mg (3 capsules) per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration:

Medicaid/HIM – 6 months



Commercial – Length of Benefit

B. NTRK Fusion-Positive Solid Tumor (must meet all):

- 1. Diagnosis of a solid tumor (see Appendix D for examples);
- 2. Prescribed by or in consultation with an oncologist;
- 3. Age \geq 12 years;
- 4. Meets one of the following (a or b):
 - a. Disease is metastatic;
 - b. Member has failed or is not a candidate for primary therapy (e.g., surgery, chemotherapy, radiation);
- 5. Tumor is positive for an NTRK gene fusion (e.g., ETV6-NTRK3, TPM3-NTRK1) without a known resistance mutation;
- 6. Member has not received prior NTRK targeted therapy (e.g., Vitrakvi®);
- 7. Request meets one of the following (a, b, or c):*
 - a. Adults: Dose does not exceed 600 mg (3 capsules) per day;
 - b. Pediatrics: Dose does not exceed any of the following (i, ii, or iii):
 - i. BSA $> 1.50 \text{ m}^2$: 600 mg PO QD;
 - ii. BSA 1.11 to 1.50 m²: 500 mg PO QD;
 - iii. BSA 0.91 to 1.10 m²: 400 mg PO QD;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

C. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

- 1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Rozlytrek for a covered indication and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, request meets one of the following (a, b, or c):*
 - a. Adults: New dose does not exceed 600 mg (3 capsules) per day;
 - b. Pediatrics: New dose does not exceed any of the following (i, ii, or iii):
 - i. BSA $> 1.50 \text{ m}^2$: 600 mg PO QD;
 - ii. BSA 1.11 to 1.50 m²: 500 mg PO QD;
 - iii. BSA 0.91 to 1.10 m²: 400 mg PO QD;
 - c. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

^{*}Prescribed regimen must be FDA-approved or recommended by NCCN.



Approval duration: Medicaid/HIM – 12 months Commercial – Length of Benefit

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration NSCLC: non-small cell lung cancer

NCCN: National Comprehensive Cancer NTRK: neurotrophic tyrosine receptor kinase

Network

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

None reported

Appendix D: Examples of Solid Tumors

(Examples are drawn from the Rozyltrek pivotal trials, as described in the FDA prescribing information, as well as the National Comprehensive Center Network (NCCN) Rozyltrek compendium.)

- Breast cancer
- Central nervous system cancers
- Cholangiocarcinoma
- Colorectal cancer
- Esophageal and esophagogastric junction cancers
- Gastric cancers
- Gynecological cancers (e.g., epithelial ovarian cancer/fallopian tube cancer/primary peritoneal cancer, uterine cancers, vulvar cancers (squamous cell), cervical cancers)
- Hepatobilliary cancers
- Lung cancer
- Neuroendocrine cancers



- Pancreatic cancer
- Salivary gland tumor
- Soft tissue sarcoma (e.g., retroperitoneal/intraabdominal, angiosarcoma, rhabdosarcoma, sarcoma of the extremity, solitary fibrous tumor, superficial trunk, undifferentiated pleomorphic sarcoma, extremity/body wall, or head/neck)
- Thyroid cancer (papillary, Hurthle cell, anaplastic, or follicular carcinoma)

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
ROS1-positive NSCLC	Adults: 600 mg PO QD	600 mg/day
NTRK fusion-positive	Adults: 600 mg PO QD	600 mg/day
solid tumor	Pediatrics (≥ 12 years of age) by body	
	surface area (BSA):	
	• BSA $> 1.50 \text{ m}^2$: 600 mg PO QD	
	• BSA 1.11 to 1.50 m ² : 500 mg PO QD	
	• BSA 0.91 to 1.10 m ² : 400 mg PO QD	

VI. Product Availability

Capsules: 100 mg, 200 mg

VII. References

- 1. Rozlytrek Prescribing Information. South San Francisco, CA: Genentech USA, Inc.; August 2019. Available at: https://www.gene.com/download/pdf/rozlytrek_prescribing.pdf. Accessed July 14, 2020.
- 2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug compendium. Accessed July 14, 2020.
- 3. National Comprehensive Cancer Network. Non-Small Cell Lung Cancer Version 6.2020. Available at: https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed July 14, 2020.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	10.01.19	11.19
Ad hoc change: NTRK fusion tumors: no known resistance	11.19.19	
mutation added for clarity, pediatric dosing details added.		
4Q 2020 annual review: no significant changes; finalized HIM line	07.14.20	11.20
of business per August SDC and prior clinical guidance; updated		
Appendix D with additional examples of solid tumors per NCCN		
Compendium; references reviewed and updated.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program



approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.



Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2019 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.