

## **Clinical Policy: Pegfilgrastim (Neulasta), Pegfilgrastim-jmdb (Fulphila), Pegfilgrastim-cbqv (Udenyca), Pegfilgrastim-bmez (Ziextenzo)**

Reference Number: IN.CP.PHAR.296

Effective Date: 12.01.16

Last Review Date: 07.21

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Pegfilgrastim (Neulasta®) and its biosimilars, pegfilgrastim-jmdb (Fulphila™), pegfilgrastim-cbqv (Udenyca™) and pegfilgrastim-bmez (Ziextenzo™), are leukocyte growth factors.

### **FDA Approved Indication(s)**

Neulasta, Fulphila, Udenyca, and Ziextenzo are indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia (FN).

Neulasta is also indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of acute radiation syndrome).

Limitation(s) of use: Neulasta, Fulphila, Udenyca, and Ziextenzo are not indicated for the mobilization of peripheral blood progenitor cells for hematopoietic stem cell transplantation.

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of Managed Health Services (MHS) that Neulasta, Fulphila, Udenyca, and Ziextenzo are **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Chemotherapy-Induced Neutropenia (must meet all):**

1. Diagnosis of non-myeloid malignancy;
2. Prescribed for use following myelosuppressive chemotherapy;
3. Request is for a biosimilar pegfilgrastim product (i.e., Fulphila, Udenyca, Ziextenzo); or member has a contraindication to a biosimilar product.
4. Dose does not exceed 6 mg (1 syringe) per chemotherapy cycle.

##### **Approval duration:**

**Medicaid** – 6 months

##### **B. Acute Radiation Syndrome (must meet all):**

1. Prescribed for use following suspected or confirmed acute exposure to myelosuppressive doses of radiation;

## CLINICAL POLICY

Pegfilgrastim, Pegfilgrastim-jmdb, Pegfilgrastim-cbqv,  
Pegfilgrastim-bmez



2. Dose does not exceed two 6 mg doses administered one week apart.

**Approval duration:**

**Medicaid** – 6 months

### **C. Bone Marrow Transplantation (off-label) (must meet all):**

1. Prescribed for one of the following (a or b):
  - a. Supportive care post autologous hematopoietic cell transplantation;
  - b. Mobilization of peripheral-blood progenitor cells prior to autologous transplantation;
2. Failure of both of the following (a and b), unless contraindicated or clinically significant adverse effects are experienced:
  - a. Neupogen<sup>®</sup>, Nivestym<sup>™</sup>, Granix<sup>®</sup>, or Zarxio<sup>®</sup>;
  - b. Leukine<sup>®</sup>;

*\*Prior authorization may be required for Neupogen, Nivestym, Granix, Zarxio, and Leukine*
3. Request meets one of the following (a or b):
  - a. Dose does not exceed 6 mg/dose;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:**

**Medicaid**– 6 months

### **D. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

## **II. Continued Therapy**

### **A. All Indications in Section I (must meet all):**

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed one of the following (a, b, or c):
  - a. Chemotherapy-induced neutropenia: 6 mg administered once per chemotherapy cycle;
  - b. Acute radiation syndrome: two 6 mg doses administered one week apart;
  - c. Bone marrow transplantation: 6 mg/dose or dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:**

**Medicaid** – 6 months

### **B. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

## CLINICAL POLICY

Pegfilgrastim, Pegfilgrastim-jmdb, Pegfilgrastim-cbqv,  
Pegfilgrastim-bmez



- Approval duration: Duration of request or 6 months (whichever is less);** or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

### III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PMN.53 for Medicaid or evidence of coverage documents.

### IV. Appendices/General Information

*Appendix A: Abbreviation/Acronym Key*

ANC: absolute neutrophil count

NCCN: National Comprehensive Cancer  
Network

FDA: Food and Drug Administration

FN: febrile neutropenia

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Neupogen (filgrastim), Zarxio (filgrastim-sndz), Granix (tbo- filgrastim), filgrastim-aafi (Nivestym)	<b>Supportive care post autologous hematopoietic cell transplantation</b> 10 mcg/kg IV or SC infusion QD  <b>Mobilization of peripheral-blood progenitor cells prior to autologous transplantation</b> 10 mcg/kg SC bolus or continuous infusion QD	10 mcg/kg/day  10 mcg/kg/day
Leukine (sargramostim)	<b>Supportive care post autologous hematopoietic cell transplantation</b> 250 mcg/m <sup>2</sup> /day IV  <b>Mobilization of peripheral-blood progenitor cells prior to autologous transplantation</b> 250 mcg/m <sup>2</sup> /day IV or SC QD	500 mcg/m <sup>2</sup> /day  250 mcg/m <sup>2</sup> /day

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): history of serious allergic reactions to human granulocyte colony-stimulating factors such as pegfilgrastim or filgrastim products
- Boxed warning(s): none reported

## CLINICAL POLICY

### Pegfilgrastim, Pegfilgrastim-jmdb, Pegfilgrastim-cbqv, Pegfilgrastim-bmez



#### Appendix D: General Information

- Neutropenia is defined as an absolute neutrophil count (ANC) of < 500 neutrophils/mcL or an ANC of < 1,000 neutrophils/mcL and a predicted decline to  $\leq$  500 neutrophils/mcL over the next 48 hours. Neutropenia can progress to FN, defined as a single temperature of  $\geq$  38.8 C orally or  $\geq$  38.0 C over 1 hour.
- The development of FN is a common dose-limiting toxicity of many chemotherapy regimens. This risk is directly related to the intensity of the chemotherapy regimen. Chemotherapy regimens that have an incidence of FN greater than 20% in clinical trials in chemotherapy naïve patients are considered by the National Comprehensive Cancer Network (NCCN) panel at high risk. Prophylaxis with myeloid growth factors is recommended at this level of risk (category 1 recommendation). NCCN Compendium recommend prophylaxis be considered in intermediate-risk (10-20% overall risk of FN) patients (category 2A recommendation). In addition to chemotherapy regimens, other risk factors such as: treatment-related, patient related, cancer-related, and co-morbidities have also been associated with an increased risk of FN. Therefore, the type of chemotherapy regimen is only one component of the risk assessment.
- Harvesting of peripheral blood stem cells prior to autologous stem-cell transplantation has a recommendation of Class IIa in DRUGDEX.
- The NCCN Compendium recommends Neulasta for supportive care post autologous hematopoietic cell transplant (category 2A).

#### V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Pegfilgrastim (Neulasta), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-bmez (Ziextenzo)	Myelosuppressive chemotherapy	6 mg administered SC once per chemotherapy cycle. Do not administer between 14 days before and 24 hours after administration of cytotoxic chemotherapy.  Weight based dosing for pediatric patients < 45 kg	6 mg/dose
Pegfilgrastim (Neulasta)	Members acutely exposed to myelosuppressive doses of radiation	Two doses, 6 mg each, administered SC one week apart. Administer the first dose as soon as possible after suspected or confirmed exposure to myelosuppressive doses of radiation, and a second dose one week after  Weight based dosing for pediatric patients < 45 kg	6 mg/dose

## CLINICAL POLICY

Pegfilgrastim, Pegfilgrastim-jmdb, Pegfilgrastim-cbqv,  
Pegfilgrastim-bmez



### VI. Product Availability

Drug Name	Availability
Pegfilgrastim (Neulasta)	<ul style="list-style-type: none"><li>• Injection: 6 mg/0.6 mL solution in a single-dose prefilled syringe for manual use only</li><li>• Injection: 6 mg/0.6 mL solution in a single-dose prefilled syringe co-packaged with the on-body injector</li></ul>
Pegfilgrastim-jmdb (Fulphila)	<ul style="list-style-type: none"><li>• Injection: 6 mg/0.6 mL solution in a single-dose prefilled syringe for manual use only</li></ul>
Pegfilgrastim-cbqv (Udenyca)	<ul style="list-style-type: none"><li>• Injection: 6 mg/0.6 mL solution in a single-dose prefilled syringe for manual use only</li></ul>
Pegfilgrastim-bmez (Ziextenzo)	<ul style="list-style-type: none"><li>• Injection: 6 mg/0.6 mL solution in a single-dose prefilled syringe for manual use only</li></ul>

### VII. References

1. Neulasta Prescribing Information. Thousand Oaks, CA: Amgen, Inc.; April 2019. Available at [www.neulasta.com](http://www.neulasta.com). Accessed May 15, 2019.
2. National Comprehensive Cancer Network. Myeloid Growth Factors Version 2.2019. Available at: [http://www.nccn.org/professionals/physician\\_gls/pdf/myeloid\\_growth.pdf](http://www.nccn.org/professionals/physician_gls/pdf/myeloid_growth.pdf). Accessed May 15, 2019.
3. Pegfilgrastim. In: National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at NCCN.org. Accessed May 15, 2019.
4. DRUGDEX<sup>®</sup> System [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed May 15, 2019.
5. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2018. Available at: <http://www.clinicalpharmacology-ip.com/>. Accessed May 15, 2019.
6. Fulphila Prescribing Information. Zurich, Switzerland: Mylan GmbH; March 2019. Available at: [www.fulphila.com](http://www.fulphila.com). Accessed May 15, 2019.
7. Udenyca Prescribing Information. Redwood City, CA: Coherus BioSciences, Inc.; April 2019. Available at: [www.udenyca.com](http://www.udenyca.com). Accessed May 15, 2019.
8. Ziextenzo Prescribing Information. Princeton, NJ: Sandoz Inc.; November 2019. Available at: [www.ziextenzo.com](http://www.ziextenzo.com). Accessed February 12, 2020.

### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J2505	Injection, pegfilgrastim, 6 mg
Q5111	Injection, pegfilgrastim-cbqv, biosimilar, (Udenyca), 0.5 mg

## CLINICAL POLICY

Pegfilgrastim, Pegfilgrastim-jmdb, Pegfilgrastim-cbqv,  
Pegfilgrastim-bmez



HCPCS Codes	Description
Q5108	Injection, pegfilgrastim-jmdb, biosimilar, (Fulphila), 0.5 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Neulasta is split from CP.PHAR.26.Colony Stimulating Factors 2015, and converted to a new template. Max dose and contraindications added per PI. Labeled use: Acute radiation syndrome added per PI. Removed “Neulasta will not be given from 14 days before to 24 hours after chemotherapy.” Off-label use: Posttransplant support added per NCCN.	11.16	12.16
Updated template and references. Removed off-label use.	08.16.17	11.17
3Q 2018 annual review: policies combined for Medicaid and Commercial lines of business; added HIM line of business; added off-label indications for mobilization of peripheral-blood progenitor cells and supportive care post autologous hematopoietic cell transplantation with redirection to FDA approved treatments Leukine and Neupogen, Granix, or Zarxio; references reviewed and updated.	05.08.18	08.18
Newly FDA-approved biosimilar added: Fulphila; references reviewed and updated.	07.31.18	02.19
Newly FDA-approved biosimilar added: Udenyca.	03.21.19	
3Q 2019 annual review: added Nivestym to list of filgrastim products required for bone marrow transplant indication, updated HCPCS coding table to include biosimilar products; references reviewed and updated.	07.17.20	07.19
Created Local Policy to meet state requirements: Removed redirection to Zarxio	10.08.19	
RT4: added Ziextenzo to policy.	02.12.20	
Added redirection to a biosimilar pegfilgrastim	08.27.20	
Annual review – No Changes		07.21

### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health

## CLINICAL POLICY

### Pegfilgrastim, Pegfilgrastim-jmdb, Pegfilgrastim-cbqv, Pegfilgrastim-bmez



plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note: For Medicaid members,** when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.