

Clinical Policy: Intestinal and Multivisceral Transplant

Reference Number: CP.MP.58

Last Review Date: 06/19

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Medical necessity criteria for the review of intestinal and multivisceral transplant requests

Policy/Criteria

It is the policy of health plans affiliated with Centene Corporation® that any of the intestinal and/or multivisceral transplantation procedures listed in **I** are **medically necessary** for pediatric and adult members to restore function in those with irreversible intestinal failure when meeting the criteria in section **II**:

I. Transplantation Procedures

- A. Isolated *intestinal transplantation* is indicated for members who have only isolated intestinal failure and no liver disease.
- B. Combined *intestinal and liver transplant* is indicated in those with intestinal failure and end stage liver disease.
- C. *Multivisceral transplant* is indicated in those with intestinal failure and the presence of neuropathy or extensive mesenteric thrombosis.

II. Procedure Criteria: Members must have one of the indications in **A** and none of the contraindications in **B**:

- A. Indications, any one of the following:
 - 1. Failure of total parenteral nutrition (TPN) as indicated by one of the following:
 - a. Impending or overt liver failure due to TPN, indicated by elevated serum bilirubin and/or liver enzymes, splenomegaly, thrombocytopenia, gastro-esophageal varices, coagulopathy, peristomal bleeding, or hepatic fibrosis/cirrhosis;
 - b. Thrombosis of ≥ 2 central veins, including jugular, subclavian, and femoral veins;
 - c. Two or more episodes of systemic sepsis due to line infection, per year, or one episode of septic shock, acute respiratory distress syndrome, and/or line related fungemia;
 - d. Frequent episodes of dehydration despite IV fluid supplementation;
 - e. Other complications leading to loss of vascular access;
 - 2. High risk of death if transplant is not performed;
 - 3. Severe short bowel syndrome (gastrostomy, duodenostomy, and/or residual small bowel <10 cm in infants and <20 cm in adults);
 - 4. Frequent hospitalizations for complications directly related to intestinal failure;
 - 5. Significant hepatic cirrhosis associated with diffuse post-mesenteric thrombosis;
- B. Does not have ANY of the following contraindications:
 - 1. Malignancy in the past two years, except for non-melanoma localized skin cancer that has been treated appropriately;
 - 2. Untreatable significant dysfunction of another major organ system, unless combined organ transplantation can be performed;

CLINICAL POLICY

Intestinal and Multivisceral Transplant

3. Presence of other GI diseases;
4. Acute medical instability, including, but not limited to, acute sepsis or myocardial infarction;
5. Uncorrectable bleeding diathesis;
6. Chronic infection with highly virulent and/or resistant microbes that are poorly controlled pre-transplant;
7. Current non-adherence to medical therapy or a history of repeated or prolonged episodes of non-adherence to medical therapy that are perceived to increase the risk of non-adherence after transplantation;
8. Psychiatric or psychological condition associated with the inability to cooperate or comply with medical therapy;
9. Absence of an adequate or reliable social support system;
10. Severely limited functional status with poor rehabilitation potential;
11. Substance abuse or dependence (including tobacco and alcohol) without convincing evidence of risk reduction behaviors, such as meaningful and/or long-term participation in therapy for substance abuse and/or dependence. Serial blood and urine testing may be used to verify abstinence from substances of concern.

Background

Intestinal transplantation is a therapeutic option for patients with intestinal failure. Intestinal failure is the loss of absorptive capacity of the small bowel secondary to severe primary gastrointestinal disease or surgically induced short bowel syndrome (SBS). The normal small intestine length varies widely, ranging from 3 to 8 meters. SBS occurs when there is approximately < 200 cm of small bowel remaining.

Multi-visceral transplantation includes the stomach, duodenum, pancreas, liver, and small intestine. A modified version excludes the liver if the recipient's liver is normal. A kidney transplant is occasionally included if the recipient has end-stage renal disease.⁵

Common indications for intestinal transplantation in children include:

- Small bowel atresia
- Gastroschisis
- Aganglionosis (Hirschsprung's disease)
- Infections such as necrotizing enterocolitis and mesenteric ischemia
- Intestinal pseudo-obstruction
- Microvillus inclusion disease
- Short gut syndrome
- Trauma
- Crohn's disease
- Midgut volvulus
- Massive resection secondary to tumor

Common indications for intestinal transplantation in adults include:

- Short gut syndrome
- Mesenteric ischemia following thrombosis, embolism, volvulus, or trauma
- Crohn's disease
- Small bowel tumors
- Small bowel secretory disorders
- Tumors of mesenteric root and retroperitoneum
- Trauma
- Volvulus
- Pseudo-obstruction
- Radiation enteritis

CLINICAL POLICY

Intestinal and Multivisceral Transplant

Guideline Recommendations

The British Society of Gastroenterology (2006) recommends: patients with SBS, including irreversible intestinal failure, expected to die prematurely on TPN, should be referred for consideration of short bowel transplant where appropriate.

The American Society of Transplantation (AST, 2001) issued a position paper on indications for pediatric intestinal transplantation. The AST recommends intestinal transplantation only for TPN-dependent children with intestinal failure who have or are likely to develop life-threatening TPN-related complications such as liver disease, recurrent sepsis, and threatened loss of central venous access. The AST stated that intestinal transplantation should not be performed solely because of continued dependence on TPN.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2019, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
44135	Intestinal allotransplantation; from cadaver donor
44136	Intestinal allotransplantation; from living donor
44715	Backbench standard preparation of cadaver or living donor intestine allograft prior to transplantation, including mobilization and fashioning of the superior mesenteric artery and vein
44720	Backbench reconstruction of cadaver or living donor intestine allograft prior to transplantation; venous anastomosis, each
44721	Backbench reconstruction of cadaver or living donor intestine allograft prior to transplantation; arterial anastomosis, each
47135	Liver allotransplantation, orthotopic, partial or whole, from cadaver or living donor, any age
47143	Backbench standard preparation of cadaver donor whole liver graft prior to allotransplantation, including cholecystectomy, if necessary, and dissection and removal of surrounding soft tissues to prepare the vena cava, portal vein, hepatic artery, and common bile duct for implantation; without trisegment or lobe split
47144	Backbench standard preparation of cadaver donor whole liver graft prior to allotransplantation, including cholecystectomy, if necessary, and dissection and removal of surrounding soft tissues to prepare the vena cava, portal vein, hepatic artery, and common bile duct for implantation; with trisegment split of whole liver graft into two partial liver grafts (i.e., left lateral segment (segments II and III) and right trisegment (segments I and IV through VIII))

CLINICAL POLICY
Intestinal and Multivisceral Transplant

CPT® Codes	Description
47145	Backbench standard preparation of cadaver donor whole liver graft prior to allotransplantation, including cholecystectomy, if necessary, and dissection and removal of surrounding soft tissues to prepare the vena cava, portal vein, hepatic artery, and common bile duct for implantation; with lobe split of whole liver graft into two partial liver grafts (i.e., left lobe (segments II, III, and IV) and right lobe (segments I and V through VIII))
47146	Backbench reconstruction of cadaver or living donor liver graft prior to allotransplantation; venous anastomosis, each
47147	Backbench reconstruction of cadaver or living donor liver graft prior to allotransplantation; arterial anastomosis, each

HCPCS Codes	Description
S2053	Transplantation of small intestine and liver allografts
S2054	Transplantation of multivisceral organs
S2055	Harvesting of donor multivisceral organs, with preparation and maintenance of allografts; from cadaver donor
S2152	Solid organ(s), complete or segmental, single organ or combination of organs; deceased or living donor(s), procurement, transplantation, and related complications; including: drugs; supplies; hospitalization with outpatient follow-up; medical/surgical, diagnostic, emergency, and rehabilitative services; and the number of days of pre- and post-transplant care in the global definition

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
A41.89	Other specified sepsis
A41.9	Sepsis, unspecified organism
K50.00-K52.9	Non-infective colitis and enteritis
K55.011-K55.9	Vascular disorders of intestine
K56.0-K56.7	Paralytic ileus and intestinal obstruction without hernia
K70.0-K77	Diseases of liver
P76.8	Other specified intestinal obstruction of newborn
P77.1-P77.9	Necrotizing enterocolitis of newborn
Q41.0-Q41.9	Congenital absence, atresia and stenosis of small intestine
R65.20-R65.21	Severe sepsis
S35.299(A/D/S)	Unspecified injury of branches of celiac and mesenteric artery, initial, subsequent encounter and sequela
T86.850-T86.859	Complication of intestine transplant
Z94.82	Intestine transplant status

Reviews, Revisions, and Approvals	Date	Approval Date
Policy developed Specialist review (Surgical Transplant)	02/14	02/14
References reviewed and updated Formatting and template updated	02/15	02/15
Minor language updates for clarification References reviewed and updated Formatting and template updated	02/16	02/16
Consolidated criteria from HN policy. Edited contraindications to be more consistent across transplant policies: Changed substance abuse to substance abuse or dependence, and added option for blood/urine testing if needed; added bleeding diatheses; reworded other contraindications for clarity. Added ICD-10 Codes. Added additional CPT and HCPCS codes.	8/16	09/16
References reviewed and updated. Some re-wording for clarity.	09/17	09/17
References reviewed and updated.	06/18	06/18
References reviewed and updated. Added CPT-47135	06/19	06/19

References

1. American Gastroenterological Association Clinical Practice Committee. AGA technical review on short bowel syndrome and intestinal transplantation. *Gastroenterology* 2003;124:1111-1134.
2. Bischel MD. Medical review criteria guidelines for managed care: Intestinal/multivisceral transplants. Apollo Managed Care Inc. Twelfth Edition, 2013.
3. Buchman, Alan L., et al. "Parenteral Nutrition-Associated Liver Disease and the Role for Isolated Intestine and Intestine/Liver Transplantation." *Hepatology*, 2006;43-1: pp. 9-19., doi:10.1002/hep.20997.
4. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for intestinal and multi-visceral transplantation. Effective May 11, 2006.
5. Khan FA Selvaggi G. Overview of intestinal and multivisceral transplantation. In: UpToDate, Brown RS (Ed), UpToDate, Waltham, MA. Accessed on 05/20/19.
6. Gilroy RK. Intestinal and multivisceral transplantation. *Medscape Reference*, Shaprio R (Ed), Apr 6, 2015. Updated May 24, 2018
7. Greenstein SM. Intestinal transplantation. *Medscape Reference*, Mancini MC (Ed). Updated May 24, 2018.
8. Hayes. Health Technology Brief. Living Related Donor Small Bowel Transplantation for Intestinal Failure. Updated April 22, 2016. Archived July 26, 2018.
9. Kato T, et al. Intestinal and multivisceral transplantation in children. *Ann Surg.* 2006; 243(6):756-64; discussion 764-6.
10. Kaufman SS, Atkinson JB, Bianchi A, et al. American Society of Transplantation. Indications for pediatric intestinal transplantation: a position paper of the American Society of Transplantation. *Pediatr Transplant.* 2001;5(2):80-87.
11. Kubal CA, Mangus RS, Tector AJ. Intestine and multivisceral transplantation: Current status and future directions. *Curr Gastroenterol Rep.* 2015;17(1):427

CLINICAL POLICY**Intestinal and Multivisceral Transplant**

12. Levitsky J, Doucette K. Viral Hepatitis in Solid Organ Transplantation. *American Journal of Transplantation*. 2013; 13: 147–168.
13. Llad L, et al. Management of portal vein thrombosis in liver transplantation: influence on morbidity and mortality. *Clinical Transplantation*. 2007;21: 716721.
14. Mangus RS, et al. Multivisceral transplantation: expanding indications and improving outcomes. *J Gastrointest Surg*. 2013 Jan;17(1):179-86.
15. Nightingale J, Woodward JM. Small Bowel and Nutrition Committee of the British Society of Gastroenterology. Guidelines for management of patients with a short bowel. *Gut*. 2006;55(Suppl IV):iv1-iv12.
16. Nishida S. Pediatric intestinal and multivisceral transplantation. Medscape Reference, Greenstein SM (Ed), Jan 06, 2016. <http://emedicine.medscape.com/article/1013915-overview>
17. Troppmann C, Gruessner RW. Intestinal transplantation. In: *Surgical Treatment: Evidence-Based and Problem-Oriented*, Holzheimer RG, Mannick JA (Ed), Munich: Zuckschwerdt; 2001.
18. Tzakis AG, et al. 100 multivisceral transplants at a single center. *Ann Surg*. 2005 October; 242(4): 480–493.
19. U.S. Department of Veteran Affairs. HIV/ AIDS. Laboratory Tests and HIV: Entire Lesson. Available at: <http://www.hiv.va.gov/patient/diagnosis/labtests-single-page.asp>
20. Vianna RM, et al. Multivisceral transplantation for diffuse portomesenteric thrombosis. *Ann Surg*. 2012 Jun;255(6):1144-50.
21. Wu G, Cruz RJ. Liver inclusion improves outcomes of intestinal retransplantation in adults. *Transplantation* 2015; 99:1265.
22. Dunn CP. Intestinal Transplantation. Medscape Reference. Mancini MC (Ed), January 18, 2017.
23. Bharadwaj S, Tandon P, Gohel TD, et al. Current status of intestinal and multivisceral transplantation. *Gastroenterol Rep (Oxf)*. 2017 Feb;5(1):20-28.
24. Pironi L, Arends J, Bozzetti F, et al. ESPEN guidelines on chronic intestinal failure in adults. *Clin Nutr*. 2016 Apr;35(2):247-307
25. Hawksworth JS, Desaid CS, Khan KM, et al. Visceral transplantation in patients with intestine-failure associated liver disease: evolving indications, graft selection, and outcomes. *Am J Transplant*. 2018 Jun; 18(6): 1312–1320.
26. Camilleri M. Chronic intestinal pseudo-obstruction. In: UpToDate, Friedman LS (Ed.), Waltham, MA. Accessed May 20, 2019.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health

CLINICAL POLICY

Intestinal and Multivisceral Transplant



plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

Note: For Medicare members, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

CLINICAL POLICY
Intestinal and Multivisceral Transplant



©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.